


1



2 **Disclaimer:**

3

4 As a condition to the use of this document and the information contained herein, the
5 Facial Identification Scientific Working Group (FISWG) requests notification by e-mail
6 before or contemporaneously to the introduction of this document, or any portion
7 thereof, as a marked exhibit offered for or moved into evidence in any judicial,
8 administrative, legislative, or adjudicatory hearing or other proceeding (including
9 discovery proceedings) in the United States or any foreign country. Such notification
10 shall include: 1) the formal name of the proceeding, including docket number or similar
11 identifier; 2) the name and location of the body conducting the hearing or proceeding;
12 and 3) the name, mailing address (if available) and contact information of the party
13 offering or moving the document into evidence. Subsequent to the use of this document
14 in a formal proceeding, it is requested that FISWG be notified as to its use and the
15 outcome of the proceeding. Notifications should be sent to: chair@fiswg.org

16 **Redistribution Policy:**

17

18 FISWG grants permission for redistribution and use of all publicly posted documents
19 created by FISWG, provided that the following conditions are met:

20

21 Redistributions of documents, or parts of documents, must retain the FISWG cover
22 page containing the disclaimer.

23


24 Neither the name of FISWG, nor the names of its contributors, may be used to endorse
25 or promote products derived from its documents.

26

27 Any reference or quote from a FISWG document must include the version number (or
28 creation date) of the document and mention if the document is in a draft status.

29

30



Physical Stability of Facial Features of Adults

31 Purpose

32 The purpose of this document is to describe the relative physical stability of facial
33 features of adults when assessing the observed component characteristics within a
34 single living subject as an aid to the facial comparison practitioner.

35 Scope

36 **This document is meant to be used in conjunction with [ASTM E3149-18] for**
37 **Morphological Analysis**

38 This document refers only to images appearing to be adult (i.e., post-pubescent)
39 subjects and does not address the stability of features in children due to rapid
40 developmental changes. This document does not cover the entire comparison
41 process. Other documents (standards/guidelines/best practices) should be referenced
42 regarding the effects of imaging conditions (e.g., illumination, pose, resolution) have on
43 feature appearance, comparison and evaluation strategies, and opinion scales.

44 **Limitations**

45 The stability assessments provided in this document are primarily a consensus
46 opinion of experienced practitioners informed by discussions with medical and
47 academic professionals, and the limited scientific literature available (as noted in the
48 specific feature tables). As more research is published, the stability assessments may
49 be updated.

50 **Introduction**

51 Over time, images of the same person may contain apparent differences due to
52 anticipated changes (e.g., aging and expression) or unanticipated changes (e.g.,
53 fluctuations in weight, health, or effects of substance use). Other visible differences may
54 be due to intentional alterations to appearance. While some changes, such as
55 expression and weight gain/loss, may be transient, others may result in a permanent
56 change in appearance. When conducting a forensic facial one-to-one comparison of
57 images captured of the same person at different time periods, the practitioner must
58 consider these potential variances when forming an opinion. This document is intended
59 to bring these concerns to the attention of the practitioner, ensure reliability of analysis,
60 and provide a pathway for the development of further standards in this area.

61 ***Factors Affecting Physical Stability of Facial Components***

62 The following factors affect the physical stability of facial features and their
63 components on an individual face (regardless of imaging conditions), listed in no
64 particular order: expression, aging (short and long term), marked weight change,

65 change in health, and intentional alteration. For each factor, stability is assessed for that
66 factor taken in isolation. In other words, under “Expression”, the stability is ONLY
67 assessed as a function of changes in expression. However, multiple factors can act
68 concurrently on the same components which the practitioner will have to take into
69 consideration. Not all factors will affect components at the same time or in the same
70 way and some components may not be affected at all. The physical stability of features
71 under each of these factors is presented below.

72 **Expression**

73 This factor refers to any deviation from a relaxed face. A relaxed face usually
74 includes eyes open and a closed mouth; however, there are individuals for whom the
75 relaxed face includes an open mouth. A neutral expression ((non-smiling) with both
76 eyes open normally (i.e., not wide-open), and mouth closed (unless medical condition
77 precludes it¹) is generally the standard by which controlled captured images (passport,
78 ID etc.) are collected. If an individual is depicted in two images under similar imaging
79 conditions with no change in expression or any other factor, then all features should
80 appear consistent.

81 **Time-related Changes**

82 This factor refers to facial variations that occur as a person ages over time. The
83 scope of time-related changes considers (without limitation) exposure to the elements
84 (e.g., sun, wind), dental changes, skin elasticity, hair loss, hyperpigmentation or

¹ ANSI/NIST-ITL 1-2011, August 22, 2016

85 hypopigmentation that may occur during this progression. For the purposes of this
86 document, time-related changes are discussed in two categories:

87 ▶ **Short term** refers to periods of five (5) years or less.

88 ▶ **Long term** refers to periods in excess of five (5) years.

89 The practitioner should be aware that these time frames are general and may need to
90 be adjusted as the practitioner considers the differing rates of change applicable to the
91 person's gender and ethnicity. If an individual is depicted in two contemporaneous
92 images under similar imaging conditions with no other factors changed, then all features
93 would be expected to appear consistent.

94 **Marked Weight Change**

95 This factor refers to the variations to the face that occur as a function of observable
96 weight loss or gain. The specific details of these variations will differ from person to
97 person. If an individual is depicted in two images under similar imaging conditions with a
98 negligible change in weight or other factors, then all features would be expected to
99 appear consistent.

100 **Changes in Health**

101 This factor refers to variations to the face that occur as a function of changes in
102 health. A comprehensive delineation of the effect of all potential health conditions is
103 beyond the scope of this document. If an individual is depicted in two images under similar
104 imaging conditions with no substantial change in health or other factors, then all
105 features would be expected to appear consistent.

106 *Note on Trauma, Inflammation, Tumors, and Substance Use:* Trauma to the head,
107 inflammation (e.g., due to infection or allergic reaction), or tumors can alter any facial
108 feature or portion of the face temporarily or permanently. As a result, the stability of all
109 facial features affected by trauma, inflammation, or tumors is low and will not be
110 delineated in the tables below. Additionally, substance use depends on multiple different
111 factors to include, but not limited to, the body chemistry of the person, the substance
112 being used, and the amount of use. Therefore, substance use will not be delineated in
113 the tables below.

114 **Intentional Alteration**

115 This factor refers to variations to the face that occur as a function of deliberate
116 modifications. These variations can be temporary or permanent. Changes may result
117 from visible modifications to the skin surface such as facial hair, tattoos, piercings*, or
118 cosmetics/makeup. Changes may also result from modifications below the skin surface
119 due to cosmetic, dental, or reconstructive procedures. Modifications due to cultural
120 practices can also affect the appearance of facial features. If an individual depicted in
121 two images under similar imaging conditions with no intentional alterations or changes
122 in other factors, then all features would be expected to appear consistent. Intentional
123 alterations are component characteristics whose stability must also be considered.

124 **Note on Piercings:* The presence of piercings may have a widely variable impact on
125 most component characteristics.

126 **Stability Tables**

127 The tables below define component characteristics as having either “High”, “Medium”
 128 or “Low” stability in the same person under the factor in question. The potential for
 129 change in the characteristic descriptors will determine this stability. If a feature with low
 130 stability is relied upon in a comparison, there is a higher risk of the comparison results
 131 being incorrect. The risk of inaccuracy increases as more low stability features are
 132 relied upon.

- 133 ▶ **High stability (H)** features exhibit little to no change.
- 134 ▶ **Medium stability (M)** features may exhibit moderate changes.
- 135 ▶ **Low stability (L)** features may exhibit substantial changes.

136 **NOTE: Each table is accompanied by text describing conditions under which**
 137 **each factor affects the stability of the corresponding component characteristic.**

Skin						
Skin appearance is extremely variable. Skin appearance may be affected by emotion, hormone levels, temperature, fatigue, hydration, etc. Changes in health and intentional alterations (e.g., make-up, tanning, tattoos, skin bleaching, and other cosmetic procedures) may cause greater variation in skin appearance.						
Component Characteristic	Expression	Time-related Changes (Short Term)	Time-related Changes (Long Term)	Marked Weight Change	Changes in Health	Intentional Alterations
Overall Skin Appearance	M	M	L	M	L	L

Face/Head Outline						
<p>The shape of the cranial vault does not change substantially in adulthood under normal conditions, but weight fluctuation or subdermal implants may give the appearance of change.</p> <p>Changes in weight or expression affect the shape of the face with the latter dominated by movement of the lower jaw. The stability of the face shape over long periods of time may also depend on tooth and related bone loss. Changes in health and intentional alterations (e.g., disease, maxillofacial surgery, orthodontic procedures, and cosmetic procedures) may cause greater variation in the overall shape of the face.</p>						
Component Characteristic	Expression	Time-related Changes (Short Term)	Time-related Changes (Long Term)	Marked Weight Change	Changes in Health	Intentional Alterations
Shape of Cranial Vault	H	H	H	M	H	M
Overall Shape of Face	L	H	M	L	L	L

141

Face/Head Composition						
<p>While the proportions of the features of the face are less stable, the position of the eyes, ears and nose relative to each other remains stable under most conditions.</p> <p>Expressions can affect the proportions of the facial features with the greatest effect occurring with movement of the lower jaw and mouth. The stability of the proportions/position of features over long periods of time may also depend on tooth and related bone loss. Changes in health and intentional alterations (e.g., disease, maxillofacial surgery, orthodontic procedures, and cosmetic procedures) may cause greater variation in the proportions/position of features on the face.</p>						
Component Characteristic	Expression	Time-related Changes (Short Term)	Time-related Changes (Long Term)	Marked Weight Change	Changes in Health	Intentional Alterations
Proportions/ Position of Features on Face	L	H	M	M	L	L

142
143

144

Hair						
Both women and men may exhibit hair loss or other changes to the hairline and baldness pattern.						
Component Characteristic	Expression	Time-related Changes (Short Term)	Time-related Changes (Long Term)	Marked Weight Change	Changes in Health	Intentional Alterations
Hair	H	L	L	H	L	L
Forehead Hairline	H	L	L	H	L	L
Hairline Right Side Hairline Left Side	H	L	L	H	L	L
Cranial baldness pattern	H	L	L	H	L	L

145

146

Forehead						
<p>Both the forehead and the brow ridges are defined by the frontal bone. Forehead shape is not affected by hairline modifications, nor are brow ridges affected by eyebrow growth or recession.</p> <p>Brow ridge prominence can be altered in both men and women due to prolonged changes in hormone levels (e.g., menopause, human growth hormone) or surgical procedures.</p>						
Component Characteristic	Expression	Time-related Changes (Short Term)	Time-related Changes (Long Term)	Marked Weight Change	Changes in Health	Intentional Alterations
Forehead Shape	H	H	M	M	H	L
Brow Ridges	H	H	M	M	M	L

147

148

Eyebrow						
<p>Eyebrows have many characteristic descriptors which can be highly variable under various factors. With the exception of expression, which can change the shape, position and asymmetry of the eyebrows, most variability is related to changes in the hair details or alterations (e.g., grooming, tattoos). Some health conditions such as facial palsy (e.g., from stroke or viral condition) may create asymmetry to the appearance of the eyebrows.</p>						
Component Characteristic	Expression	Time-related Changes (Short Term)	Time-related Changes (Long Term)	Marked Weight Change	Changes in Health	Intentional Alterations
Right Eyebrow Left Eyebrow	L	H	L	H	L	L
Asymmetry between Right and Left Eyebrows	L	H	H	H	L	L

149

150

Eyes						
<p>Intercanthal distance does not change with the exception of trauma. Some component characteristics of the eye are affected by expression while others are not. Changes to the sclera can occur over short periods of time due to causes such as exposure to sun, wind, and other irritants. Time-related changes over the long term and marked weight change primarily affect the soft tissues. Examples include eyelid drooping, orbital fat variations, and corneal clouding. Changes in health can affect all other component characteristics of the eyes, and some conditions such as facial palsy (e.g., from stroke or viral condition) may create asymmetry to the appearance of the eyes. Intentional alterations to the eye include contact lenses, cosmetics, cosmetic procedures, tattoos, piercings, prostheses, etc.ⁱ</p>						
Component Characteristic	Expression	Time-related Changes (Short Term)	Time-related Changes (Long Term)	Marked Weight Change	Changes in Health	Intentional Alterations
Intercanthal Distance	H	H	H	H	H	H
Interpupillary Distance (IPD)	M	H	H	H	M	L
Right Eye Fissure Opening (Outline) Left Eye Fissure Opening (Outline)	L	H	M	M	L	M

Right Upper Eyelid (including lashes) Left Upper Eyelid (including lashes)	L	H	M	M	L	L
Right Lower Eyelid (including lashes) Left Lower Eyelid (including lashes)	L	H	L	M	L	L
Right Eyeball Prominence Left Eyeball Prominence	H	H	H	M	L	M
Right Eye Sclera Left Eye Sclera	H	M	M	H	L	L
Right Iris Left Iris	L	H	M	H	L	L
Right Eye Medial Canthus Left Eye Medial Canthus	H	H	H	H	L	M

Right Eye Lateral Canthus	H	H	M	H	L	M
Left Eye Lateral Canthus						
Asymmetry Between Right and Left Eyes	H	H	H	H	L	M

151

Cheeks						
<p>The apparent prominence of the cheekbones varies in relation to changes in weight, health (e.g., stroke, facial palsy, dental changes) and intentional alteration (e.g., cosmetics or cosmetic procedures). With aging (senescence) the cheekbone may appear more prominent due to decreased soft tissue and muscle mass.</p> <p>The cheek is a flexible soft tissue structure which can be affected by all of the factors. The buccal fat pad moves inferiorly down the cheek during middle-age and this creates a flatter cheek shape with less prominent cheekbones. Intentional alterations include cosmetics, surgical implants and fillers.</p>						
Component Characteristic	Expression	Time-related Changes (Short Term)	Time-related Changes (Long Term)	Marked Weight Change	Changes in Health	Intentional Alterations

Right Cheekbone	H	H	M	L	L	L
Left Cheekbone						
Right Cheek Shape (soft tissue)	L	M	L	L	L	L
Left Cheek Shape (soft tissue)						

152

Nose							
<p>The nose is a stable feature over the short term under normal conditions. All characteristic components, except the nasal root, can change with expression, and all components can be changed with intentional alterations (e.g., piercings, surgery). Over the long term, the soft tissues (including the cartilage) of the nose change in length and shape. Under marked weight changes, only the root and columella remain stable, and under health changes, only the root and body remain stable. The other characteristic components of the nose can be affected by disease (e.g., leprosy, gout), viral conditions (e.g., colds, sinusitis), and growths (e.g., polyps). The attachment of the alae to the upper lip does not change with the exception of surgery, trauma, or tumor.ⁱⁱ</p>							
Component	Characteristic	Expression	Time-related Changes (Short Term)	Time-related Changes (Long Term)	Marked Weight Change	Changes in Health	Intentional Alterations
Nasal Outline (Profile/Front view)		L	H	M	M	M	L

Nasal Root (bridge)	H	H	H	H	H	L
Nasal Body	M	H	H	M	H	L
Nasal Tip	L	H	L	M	M	L
Nasal Base	L	H	M	M	M	L
Nasal Base: Alae (Wings of nose)	L	H	M	M	M	L
Nasal Base: Nostrils (Nasal Openings)	L	H	M	M	M	L
Nasal Base: Columella (Soft tissue between Nostrils)	M	H	M	H	M	L

153

154

Ears						
<p>The ear is a very stable feature. Ear position alone can change with expression, but the ear configuration remains stable. Over the long term, the ear changes in proportions, although lengthening of the lobule dominates these changes. Marked weight changes can affect the prominence and protrusion of the ear, as well as the fat content of the lobe. Other than trauma (e.g., cauliflower ear), inflammation and tumors, health changes rarely affect the characteristic components of the ears, except in cases of unusual disease (e.g., leprosy or cysts).ⁱⁱⁱ</p>						
Component Characteristic	Expression	Time-related Changes (Short Term)	Time-related Changes (Long Term)	Marked Weight Change	Changes in Health	Intentional Alterations
Asymmetry Between Left and Right Ears	M	H	H	H	H	L
Right Ear Protrusion Left Ear Protrusion	M	H	M	L	H	L
Overall Right Ear Overall Left Ear	M	H	M	M	H	L

Right Ear Helix- Superior, Inferior (tail)	H	H	H	H	H	L
Left Ear Helix- Superior, Inferior (tail)						
Right Ear Tubercles (Auricular Tubercle)	H	H	H	H	H	L
Left Ear Tubercles (Auricular Tubercle)						
Right Ear Antihelix Left Ear Antihelix	H	H	H	H	H	L
Right Ear Crura of Antihelix (Superior, Inferior)	H	H	H	H	H	M
Left Ear Crura of Antihelix (Superior, Inferior)						
Right Ear Triangular fossa Left Ear Triangular fossa	H	H	H	H	H	M

Right Ear Crus of Helix	H	H	H	H	H	M
Left Ear Crus of Helix						
Right Ear Scaphoid Fossa	H	H	H	H	H	M
Left Ear Scaphoid Fossa						
Right Ear Concha (Superior, Inferior)	H	H	H	H	H	M
Left Ear Concha (Superior, Inferior)						
Right Ear Tragus	H	H	H	H	H	L
Left Ear Tragus						
Right Ear Antitragus	H	H	H	H	H	L
Left Ear Antitragus						
Right Ear Intertragic/ Intertragal Notch	H	H	H	H	H	M
Left Ear Intertragic/ Intertragal Notch						

Right Ear Anterior Knob	H	H	H	H	H	L
Left Ear Anterior Knob						
Right Ear Anterior Notch	H	H	H	H	H	M
Left Ear Anterior Notch						
Right Ear Posterior Auricular Furrow	H	H	H	H	H	M
Left Ear Posterior Auricular Furrow						
Right Ear Lobule (Lobe)	H	H	M	M	H	L
Left Ear Lobule (Lobe)						
Ear Abnormalities	H	H	M	H	H	L

155

156

Mouth						
<p>The mouth is the facial feature that changes the most under expression. Over a short period of time, the component characteristics of the mouth are stable, with the exception of tooth loss and tooth color (e.g., coffee stains). Over the long term the lips become thinner and the position of the mouth fissure may change due to this and any dental changes. Health changes will affect the component characteristics of the mouth in a variety of ways, such as lip shape (e.g., herpes simplex virus), asymmetry (e.g., stroke and palsy), lip creases (e.g., dehydration) or lip tone. Intentional alterations include tattoos, piercings, fillers and cosmetics. Mouth abnormalities, such as cleft lip and palate, are frequently corrected through cosmetic procedures which results in a different appearance.^{iv}</p>						
Component Characteristic	Expression	Time-related Changes (Short Term)	Time-related Changes (Long Term)	Marked Weight Change	Changes in Health	Intentional Alterations
Philtrum	L	H	L	M	M	L
Overall Mouth	L	H	L	M	M	L
Upper Lip	L	H	L	M	M	L
Lower Lip	L	H	L	M	M	L

Lip Fissure (Opening between lips)	L	M	L	M	M	L
Mouth Asymmetry	L	H	H	H	M	L
Overall Dental Occlusion (Contact between Upper and Lower Teeth)	L	M	M	H	M	L
Gnathism (apparent convexity or concavity of the mouth complex, related to the relative projection of the upper and/or lower teeth)	H	H	M	M	M	L
Characteristic Detail of Teeth	H	H	M	H	M	L
Mouth Abnormalities	H	H	H	H	M	L

158

159

Chin/Jawline						
<p>The chin, jawline and gonial angle are stable features and do not change over the short term. Long term changes relate to tooth and related bone loss, sagging due to loss of skin elasticity and changes in subcutaneous fat distribution with age. Marked weight change will alter the appearance of these features due to subcutaneous fat changes and the effects of gravity on the soft tissues. Health changes, such as hormone levels (e.g., menopause or steroid treatment), behavior (e.g., tooth grinding), disease (e.g., sialosis) and viral conditions (e.g., mumps, mononucleosis) can also affect them. Intentional alterations can include maxillofacial surgery, orthodontic treatment, cosmetic procedures, and facial hair.</p>						
Component Characteristic	Expression	Time-related Changes (Short Term)	Time-related Changes (Long Term)	Marked Weight Change	Changes in Health	Intentional Alterations
Chin (Profile and Frontal view)	M	H	M	L	M	L
Jawline (from Chin to Gonial Angle)	H	H	M	L	M	L
Gonial Angle (Angle of the jaw)	H	H	M	L	M	L

160

161

Neck						
<p>The neck is a stable feature in the short term. Long term there may be changes in muscle mass associated with aging, exercise and hormone levels, and positional changes related to posture.</p> <p>The neck will change shape in relation to expression and marked weight change. Health changes may affect the neck in relation to asymmetry (e.g., torticollis), width (e.g., goiter) and position (e.g., arthritis). Intentional alterations include body building, spinal surgery and postural alteration.</p> <p>The laryngeal prominence is a stable feature under normal conditions. The laryngeal prominence will change position in relation to expression, posture, and marked weight change will affect the apparent prominence. Health changes may affect the laryngeal prominence (e.g., goiter, mononucleosis) and intentional alterations include hormonal treatment, facial hair and cosmetic procedures.</p>						
Component Characteristic	Expression	Time-related Changes (Short Term)	Time-related Changes (Long Term)	Marked Weight Change	Changes in Health	Intentional Alterations
Neck (Overall)	M	H	L	L	L	L
Laryngeal Prominence (Adam's Apple)	M	H	H	L	M	L

162

163

Facial Hair						
<p>Following post-pubescent growth, the distribution, symmetry, and density of facial hair remain stable in all areas of the face in the short term, however the other characteristic descriptors are highly variable. Long term (e.g., aging, hormone levels) and health changes (e.g., disease, stress) are exhibited for all characteristic descriptors. Expression will change apparent facial hair position at the upper and lower lip but remain stable on the sides and neck. Marked weight change will affect the apparent position of facial hair. Intentional alterations include grooming, cosmetic procedures, hormonal treatment, and prostheses.</p>						
Component Characteristic	Expression	Time- related Changes (Short Term)	Time- related Changes (Long Term)	Marked Weight Change	Change s in Health	Intentional Alterations
Facial Hair Above Upper Lip	L	H	L	M	L	L
Facial Hair Below Lower Lip						
Facial Hair on Right Side	M	H	L	M	L	L
Facial Hair on Left Side						

Facial Hair on Neck, below Chin/Jawline	H	H	L	M	L	L
-----------------------------------------------	---	---	---	---	---	---

164

Facial Lines						
<p>All facial creases become more defined over time and the number of wrinkles will increase over time. Wrinkles are a skin aging response in relation to muscle action and decreased skin elasticity and will align perpendicular to the muscle fiber action. Some creases are related to anatomical structure, such as a bifid nasal tip, cleft chin and nasolabial folds. Creases and wrinkles are stable under normal conditions, although their prominence can be changed by some factors. Expression will make most facial lines more defined and may change their appearance and relative position, although some structural creases (e.g., bifid nasal tip and cleft chin) are unaffected by expression. Long term changes are related to intrinsic aging, lifestyle (e.g., smoking, drug use, alcohol consumption), stress, sun exposure and dehydration. Marked weight change will affect the position, definition and shape of crease patterns. Health changes (e.g., stroke, palsy) will affect creases and wrinkles in relation to asymmetry, definition, shape and number, and some conditions will obscure crease patterns (e.g., leprosy, goiter, dermatological conditions). Intentional alterations include cosmetic procedures, surgery and cosmetics.</p>						
Component Characteristic	Expression	Time- related Changes (Short Term)	Time- related Changes (Long Term)	Marked Weight Change	Change s in Health	Intentional Alterations

Frontal Lines (Forehead Wrinkles)	L	H	M	M	L	L
Vertical Glabellar Line(s)	L	M	M	M	L	L
Nasion Creases	L	H	M	M	L	L
Right Lateral Nasal Lines Left Lateral Nasal Lines	L	H	M	H	M	L
Bifid Nose Crease	H	H	H	M	M	L
Periorbital Lines Right Eye (Crow's Feet/wrinkles) Periorbital Lines Left Eye (Crow's Feet/wrinkles)	L	M	M	M	L	L
Right Superior Palpebral Crease Left Superior Palpebral Crease	L	H	M	M	L	L

(Crease between the Upper Eyelid and the Top of the Bony Orbit)						
Right Inferior Palpebral Crease Left Inferior Palpebral Crease (Crease between the Lower Eyelid and the Bottom of the Bony Orbit)	L	H	M	M	L	L
Right Infraorbital Creases Left Infraorbital Creases (Creases below the eyes)	L	H	M	M	L	L
Upper Circumoral Striae (Lip Creases) Lower Circumoral Striae (Lip Creases)	L	H	M	M	L	L

Mentolabial sulcus (Horizontal Crease or Fold between Lower Lip and Chin)	L	H	M	M	L	L
Right Nasolabial Crease/Folds Left Nasolabial Crease/Folds (Creases or Folds extending from Nose to Corners of Mouth)	L	H	M	M	L	M
Right Marionette Lines Left Marionette Lines	L	H	M	M	L	L
Cleft Chin	H	H	H	M	M	L
Right Buccal Creases/folds Left Buccal Creases/folds (cheek to chin)	L	H	M	M	L	L

Wrinkles on Neck	M	H	L	M	L	L
Other Creases	See below*					

165

166

167

*The stability of other creases depends on their location and orientation as such the stability of a specific crease cannot be generalized in this document.

Scars						
Over time scars may change in relation to visibility but are unaltered by marked weight change or health changes. Expression may change a scar in relation to apparent position and shape. Intentional alterations to conceal scars include cosmetics, facial hair, and tattoos.						
Component Characteristic	Expression	Time-related Changes (Short Term)	Time-related Changes (Long Term)	Marked Weight Change	Changes in Health	Intentional Alterations
Scars	M	M*	M	H	H	L

168

169

170

171

172

173

*Scars are extremely unstable during the scar maturation phase which can last up to one year once the wound has healed. Following that they are relatively stable features in the short term, under normal conditions.

Facial Marks						
<p>Facial marks are features that may be transient or permanent. Transient marks (e.g., acne or blemishes) are unstable, but other marks (e.g., moles or skin tags) may be stable both short and long term. Expression may change a facial mark in relation to apparent position and shape, depending on its original position, and marks on the lower face (e.g., mouth and lower jaw) will be affected more than those on the upper face (e.g., forehead and nose) by expression. Marks on the ear are unaffected by expression. Health changes (e.g., dermatological conditions, high blood pressure, sun damage) will affect the distribution, number, definition and position of some skin marks, such as freckles, blemishes, or warts. Intentional alteration includes cosmetics, surgery, facial hair, and tattoos.</p>						
Component Characteristic	Expression	Time-related Changes (Short Term)	Time-related Changes (Long Term)	Marked Weight Change	Changes in Health	Intentional Alterations
Skin Marks	M	L	L	M	L	L

174

175

Alterations

This section refers to the stability of each existing intentional alteration in isolation.

Piercings may be unstable over time due to healing and stretching, but the location relative to the pierced feature remains stable even with expression. Weight change may affect the appearance of a piercing. Intentional alterations of piercings include surgery, cosmetics, stretching and additional piercings.

Tattoos are stable in the short term but may fade or become blurred over time. Expression may alter the tattoo shape due to skin movement and marked weight change may stretch or crease a tattoo. Health changes may affect tattoos in relation to skin changes (e.g., dermatological conditions). Intentional alterations or concealment of tattoos include laser removal, cosmetics, additional tattooing or facial hair.

Because makeup is a transient alteration that is unstable over time and is not particularly affected by expression, weight changes, or changes in health, its stability relative to those factors is not addressed in the table below.

Other alterations include but are not limited to surgery, implants, and fillers.

Component Characteristic	Expression	Time-related Changes (Short Term)	Time-related Changes (Long Term)	Marked Weight Change	Changes in Health	Intentional Alterations
Piercing	H	M	L	M	M	L
Tattoo	M	M	M	M	M	L
Other	See below*					

176

177 *The stability of other alterations depends on their location, type, and orientation as
178 such the stability of a specific alteration cannot be generalized in this document.

179 **Conclusion**

180 Individuals conducting one-to-one comparison examinations must consider the
181 stability of the facial features. Failure to consider the stability of facial features may lead
182 to incorrect conclusions. Future documents will discuss the significance of this
183 information during the evaluation process for conclusion determination.

184 **References**

ⁱ Chiarella Sforza *, Gaia Grandi, Francesca Catti, Davide G. Tommasi,

Alessandro Ugolini, Virgilio F. Ferrario: Age- and sex-related changes in the soft
tissues of the orbital region

ⁱⁱ Chiarella Sforza *, Gaia Grandi, Marcio De Menezes, Gianluca M. Tartaglia, Virgilio
F. Ferrario: Age- and sex-related changes in the normal human external nose

ⁱⁱⁱ Chiarella Sforza *, Gaia Grandi, Miriam Binelli, Davide G. Tommasi, Riccardo
Rosati, Virgilio F. Ferrario: Age- and sex-related changes in the normal human ear

^{iv} Chiarella Sforza *, Gaia Grandi, Miriam Binelli, Claudia Dolci, Marcio De Menezes,
Virgilio F. Ferrario: Age- and sex-related changes in three-dimensional lip morphology

DRAFT

FISWG documents can be found at: www.FISWG.org